

Product information

【Drug Facts】



〈Kyowa-Gifu-Japan〉

For resource saving the case is combined with the Japanese instruction written on the package case. Be sure to read the instruction carefully before use. Also, please keep it in a safe place so that you can read again, when you need it.

Class ② drugs

Antihemorrhoids
(External Preparations)

(OINTMENT)

OSIRI NON



Active ingredients

	(in 100g)	Effect
Hydrocortisone acetate	0. 5g...	Relieves symptoms such as inflammation, itching, and swelling
Lidocaine	3. 0g...	Relieve pain and itching
Zinc oxide	4. 0g...	Soothes burns, eczema, genital itching, etc.
Isopropyl methyl phenol	0. 1g...	Bactericidal ingredient
Chlorpheniramine maleate	0. 2g...	Reduce itching
Allantoin	1. 0g...	Remedies for skin
Glycyrrhetic acid	0. 3g...	Anti-inflammatory ingredient
Tocopherol acetate	3. 0g...	Preventing rough skin and itching

Indication

Relief and disinfection of pain, itching, swelling, and bleeding caused by hemorrhoids (cleft hemorrhoids) and wart hemorrhoids



Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

1. Please the medicine should not be used on the following parts:

Patients with purulent hemorrhoidal lesions.

2. Please do not use this medicine for a long time



Consult a doctor for the following

1. The following persons should consult a doctor, pharmacist, or registered salesperson before using this medicine.
 - (1) Patients undergoing medical treatment from a doctor.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Patients who have experienced allergic symptoms associated with drugs, etc.
2. If the following symptoms appear after use, it may be a side effect, so stop using it immediately and consult your doctor, pharmacist, or registered salesperson with this case.

Related area	Symptoms
Skin	Rash ▪ redness, itching, swelling
Other	Irritation, Purulence

3. When symptoms do not improve even after using this medicine for about 10 days, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

Dosage and administration

.Apply an appropriate amount to the anal area 1 to 3 times a day.

When using this product

- (1) This medicine should be given to children only under adult supervision.
- (2) This medicine must be used only for the anal area.

Precautions for storage and handling

- (1) Store in a cool place free from direct sunlight by a seal tightly.
- (2) Keep all medicine out of the reach of children.

- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)

Inactive ingredients: Contains macrogol, medium chain fatty acid triglyceride, glyceryl monostearate, sorbitan sesquioleate, and petrolatum.

Contact information for the Relief System for Side Effects

Pharmaceuticals and Medical Devices Agency

http://WWW.pmda.go.jp/kenkouhigai_cmp/index.html

Telephone: 0120-149-931 (toll-free)

Questions or comments?

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.

Customer consultation room ☎ 058-246-8771



Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.

- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.