

Product information

【Drug Facts】



＜Kyowa-Gifu-Japan＞

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class ② drugs

Cold medicine

(TABLETS)

SURUURON EX JYOU PLUS

The initial symptoms of the cold start with a runny nose, stuffy nose, followed by cough, headache, chills, fever etc.

”SURUURON EX JYOU PLUS” is comprehensive cold medicines containing antipyretic analgesics, antitussive expectorants, antihistamines, etc. so as to alleviate symptoms of such colds as soon as possible.

Active ingredient (in 9 tablets/daily dose)

Purpose

Acetaminophen 300mg.....	Antipyretic analgesic
Etenzamide 400mg.....	Antipyretic analgesic
Anhydrous Caffeine 75mg.....	Headache treatment
Dihydrocodeine phosphate 24mg	Anti-cough effect
dl-Methylephedrine hydrochloride 60mg	Bronchodilator
Bromhexine hydrochloride 12mg.....	Antitussive expectorant
Chlorpheniramine maleate 7.5mg.....	Anti-allergy effect
Licorice extrac 250mg.....	Anti-inflammatory effect
(equivalent to 1,000mg of the herbal medicine Glycyrrhiza Radix)	
LUMBRICUS dry extract 77.92mg	Antitussive expectorant
(equivalent to 600mg of the traditional medicated earthworm)	
Bellflower poiwder 600mg.....	Antitussive expectorant
Ginger root dried extract 45mg.....	Anti-inflammatory effect
(equivalent to 450mg of the herbal medicine Ginseng Radix.)	
Cinnamon powder 150mg	Anti-thirsty effect

Uses

Relaxation of various symptoms of colds (runny nose, stuffy nose, sneeze, sore throat, cough, tan, chills, fever, headache, joint pain, muscle pain)

Warnings

Do not take children under 12 years old.

[To be avoided]

(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

1. Do not take to the following person:

- (1) A person with a medical history of being allergic to this medicine or substance in it.
- (2) A person who have ever had asthma taking this medicine or other cold medicine, antipyretic analgesics.

2. While taking this medicine, you should not use in combination with the following medicines:

Other medicines for cold, antipyretic analgesics, sedatives, antitussive expectorants, and internal medicines containing antihistamine etc. (internal medicine for rhinitis, motion sickness medicine, allergy medicine etc).

3. Do not drive a motor vehicle or operate machinery after using this medicine. (Symptoms as drowsiness may occur.)
4. You should not take this medicine during breastfeeding a baby. If you want take this medicine you should avoid to breastfeed your baby.
5. Do not drink alcohol before or after taking this medicine.
6. Do not use in long-term continuously.

[Consultation]

1. The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.
 - (1) A person undergoing medical treatment from a doctor or dentist..
 - (2) A pregnant woman or a woman suspected of being pregnant.
 - (3) Baby/infants/children (under 15 years old) who have varicella (chickenpox) or influenza or are suspected of having either of them.

(4) Senior persons.

(5) A person with a medical history of being allergic to medicines and other substances.

(6) A person with the following symptoms:

High fever, Swelling, Dysuria.

(7) A person diagnosed as having the following:

Thyroid dysfunction, diabetes, heart disease, hypertension, liver disease, kidney disease, gastric / duodenal ulcer, glaucoma, respiratory dysfunction, obstructive sleep apnea syndrome, obesity.

2. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, discontinue use immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

Related area	Symptoms
Skin	Rash · Redness, itching
Digestive organ	Nausea / vomiting, loss of appetite
Mental nervous system	Dizzy
Urology	Dysuria
Others	Excessive hypothermia

In very rare cases the following serious symptoms may occur. In this case, please get medical attention immediately.

Symptoms' name	Symptoms
Shock(Anaphylaxis)	Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.
Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis:	Hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.

Hepatic function failure	Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.
Kidney disorders	Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints) and diarrhea, etc. may occur.
Interstitial pneumonia	Shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance
Asthma	Symptoms, such as wheezing or hissing when breathing, and difficult breathing, etc. may occur.
Aplastic anemia	Symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.
Agranulocytosis	Symptoms, such as acute high fever, chills and sore throat, etc. may occur.
Respiratory depression	Constipation, Shortness of breath, breathing difficulties and so may appear.

3. The following symptoms may appear after taking this medicine, so if such symptoms persist or increase. Please stop taking it and consult a doctor, pharmacist or registered distributor with the Japanese package insert.

Constipation, Thirst, Sleepiness

4. If these symptoms are not good even after taking 5 or 6 times, please stop taking the medicine and consult your doctor, pharmacist or registered distributor with the Japanese package insert.

Directions

Please as possible within 30 minutes after meals.

Age	One dose	Number of doses per day
adult and children 15 years and over	3 tablets	3 times


children 12 years to under 15 years	2 tablets	3 times
children under 12 years	Do not use	

[When using this product]

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children only under adult supervision.
- (3) How to take out tablets. (as to a PTP product),

Please push the projection of the PTP sheet containing tablets strongly with your fingertips as shown below, break the aluminum sheet on the back side, push out and take it.

(If you mistakenly swallow the PTP sheet as it is, it leads to an unexpected accident such as stabbing into the esophagus mucosa)



Other information

- (1) Store in a cool and dry with less humidity place free from direct sunlight.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)
- (4) If you take medicine after dividing one packet, please fold the mouth of the packet and store it, please take it within 2 days.
- (5) Do not take this medicine after the expiration date.

Inactive ingredients:

Lactose, Polyvinyl alcohol, Methyl methacrylate copolymer, Mg Stearate, Macrogol, Talc, Silicon dioxide, Cross CMCNa, Hypromerose, Titanium oxide, Iron sesquioxide.

Questions or comments

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.
Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] **<Kyowa-Gifu-Japan>**

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the Revised date, the attached document is revised from time to time, so the English translation may not be the latest.