

Product information

【Drug Facts】



<Kyowa-Gifu-Japan>

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 3 drugs **Magnesium Salt' Laxative** (TABLETS)

TIFA S JYOU

Magnesium oxide collects water in the intestines and softens the stools, thereby promoting intestinal movement and natural defecation.

It is formulated with Coix seed, which has the effect of improving skin problems, and eliminates rough skin or pimples due to constipation.

Active ingredients (in 6tablets/daily dose)

Purpose

Magnesium Oxide 2000mg

Laxative ingredient

Coix seed extract 310mg.....

Improving Skin Problems

Uses

- Constipation
- Relief of the following symptoms associated with constipation.
Dull headache, Hot flashes, Rough skin, Pimples, Loss of appetite, Abdominal bloating, Intestinal abnormal fermentation, Hemorrhoids

Warnings

[To be avoided]

(Failure to observe the following may result in the current symptoms worsening or adverse reactions occurring.)

1. While taking this medicine, please do not take any of the following medicines.

Other laxatives (purgative)

[Consultation]

1. The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

(1) A person undergoing medical treatment from a doctor.

(2) A pregnant woman or a woman suspected of being pregnant.

(3) Senior persons.

(4) A person with the following symptoms:

Severe headache, nausea, vomiting.

(5) A person diagnosed as having the following:.

Kidney disease.

2. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, stop the use immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

Related area	Symptoms
Digestive organ	Severe stomach ache, nausea and vomiting
Cental nervous system	Strong sleep, weakening awareness
Circulatory organ	Dizziness , pulse slows down
Respiratory organ	Breathless
Others	Muscle weakness, thirst

3. The following symptoms may appear after taking it, so if such symptoms persist or increase. Please stop taking it and consult a doctor, pharmacist or registered distributor with the Japanese package insert.

Diarrhea.

4. If these symptoms do not improve after having used this medicine 5 or 6 days, stop taking the medicine, and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

Directions

Please take the next dose once a day with a glass of water before going to bed. However, please use the minimum amount at first, and increase or decrease little by little, while observing the condition of the body and of the bowel movement.

Age	One dose	Number of doses per day
adult and children 15 years and over	3~6 tablets	once (before going to bed)
children 11 years to under 15 years	2~4 tablets	
children 7 years to under 11 years	2~3 tablets	
children 5 years to under 7 years	1~2 tablets	

children under 5 years

Do not use

[When using this product]

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children only under adult supervision.

Other information

- (1) Store in a cool and dry with less humidity place free from direct sunlight by a seal tightly.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)
- (4) The filling in the bottle is to prevent the tablets from breaking during shipping. It will be unnecessary after opening, so please remove it.
- (5) Please do not take products that have passed the expiration date.

Inactive ingredients:

Cellulose, Croscarmellose sodium, Erythritol,
Aspartame (L-Phenylalanine compound), L-
Menthol and calcium stearate.

Questions or comments

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.
Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.

- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the Revised date, the attached document is revised from time to time, so the English translation may not be the latest.