

Product information

【Drug Facts】



<Kyowa-Gifu-Japan>

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 3 drugs

Multivitamin Preparations

(TABLETS)

VITAGOLD EX ACE

“VITAGOLD EX ACE” is a so-called “Vitamin B1,B6,B12 formulation”, besides contains vitamin E, gamma-oryzanol and calcium pantothenate, which are taken for symptoms such as “Physical poor condition by unknown catake.”

Active ingredients (in 3 tablets/daily dose)

Purpose

| | |
|---------------------------------------|-----------------------------|
| As Flusultiamine 100mg••••• | Vitamin B1 supplementation |
| Pyridoxine hydrochloride 100mg••••• | Vitamin B6 supplementation |
| Cyanocobalamine 1,500 μ g••••• | Vitamin B12 supplementation |
| Tocopherol succinate succinate 100mg• | Vitamin E supplementation |
| Calcium pantothenate 30mg••••• | Vitamin supplementation |
| Gamma-Oryzanol 10mg••••• | Vitamin supplementation |
| Nicotinic acid amide 60mg••••• | Vitamin supplementation |

Uses

- ◎ Alleviation of next symptoms.

Neuralgia, myalgia / arthralgia (backache, stiff neck, fifty shoulders, etc.), numbness of hands and feet, eye strain.

“However, if these symptoms do not improve after one month of use, please consult your doctor or pharmacist.”

- ◎ Supplementation of Vitamin B1、B6、B12 in the following cases.

At the time of physical fatigue, pregnancy or lactation, and physical poor conditions after or during illness.

Warnings

[Consultation]

1 The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

- (1) A person undergoing medical treatment from a doctor.
- (2) A person who have ever had allergic symptoms due to any medication.

2. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, stop the take immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

| Related area | Symptoms |
|-----------------|--|
| Skin | Rash · Redness、Itching |
| Digestive organ | Nausea · Vomiting, Stomatitis and Stomach discomfort |

3. The following symptoms may appear after taking it, so if such symptoms persist or increase. Please stop taking it and consult a doctor, pharmacist or registered distributor with the Japanese package insert.

Soft stools, Diarrhea, Constipation.

4. If symptoms do not improve after taking this medicine for about one month, stop taking the medicine, bring the Japanese package insert, and consult a doctor, pharmacist or registered distributor.

5. After taking this medicine, menstruation may come earlier than planned, and menstrual blood volume may increase slightly. If the bleeding lasts for a long time, take this document and consult a doctor, pharmacist or registered distributor.

Directions

Please take the following amount with water or lukewarm water after meals without biting.

| Age | One dose | Number of doses per day |
|--------------------------------------|-------------|-------------------------|
| adult and children 15 years and over | 2~3 tablets | once |
| children under 15 years | Do not use | |

[When using this product]

(1) Please follow the recommended dosage and directions.

Other information

- (1) Store in a cool and dry with less humidity place free from direct sunlight by a seal tightly.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (It may be misused or the quality may deteriorate.)
- (4) Do not touch the tablets with wet hands as may change its shape due to moisture.
- (5) The fillings in the bottle is to prevent the tablets from breaking down during shipping. It will be not necessary after opening the bottle, so please remove the fillings.
- (6) The "paper packet" in the bottle is a desiccant. Do not remove covered paper of it, and never take out, open and eat.
- (7) Please write the "Open Date" of this medicine on the inner lid of the case.
- (8) Do not take overdue products. In addition, please use within six months for quality maintenance after opening even if it is within expiration date.

Inactive ingredients:

Anhydrous silicic acid, Lactose, Calcium lactate, Cellulose, Carmellose, Magnesium stearate, Hypromellose, White sugar, Gelatin, Gum arabic, Calcium carbonate, Talc, Titanium oxide, Riboflavin, Carnauba wax.

Questions or comments

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.
Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the Revised date, the attached document is revised from time to time, so the English translation may not be the latest.