

## Product information

### 【Drug Facts】



＜Kyowa–Gifu–Japan＞

For resource saving the case is combined with the Japanese instruction written on the package case. Be sure to read the instruction carefully before use. Also, please keep it in a safe place so that you can read again, when you need it.

Class ② drugs      Antipruritic and Anti-inflammatory      ( CREAM )  
Drugs

## MOREFINE ASEMO CREAM

### Active ingredients

(in 100g)

### Effect

Prednisolone valeric acid ester acetic acid ester  
(PVA)      0.15g\*\*\*

Reduce skin irritation

Diphenhydramine hydrochloride      2.0g\*\*\*\*

Reduce the occurrence of itching

l-Menthol      3.5g\*\*\*\*\*

Give a refreshing feeling

l-Camphor      3.5g\*\*\*\*\*

Give a refreshing feeling

Crotamiton      5.0g\*\*\*\*\*

Reduce itching

Isopropyl methyl phenol      0.1g\*\*\*\*\*

Bactericidal ingredient

### Indication

Eczema, dermatitis, heat rash, rash, itching, insect bites,  
hives



### Precautions

#### ⊗ When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

#### 1. Please the medicine should not be used on the following parts:

- (1) Varicella (chickenpox)、Athlete's foot/tinea or purulent lesions
- (2) Periorbital area、Mucosa

#### 2. Please do not use extensively on the face.

#### 3. Please do not use this medicine for a long time



## Consult a doctor for the following

**1. The following persons should consult a doctor, pharmacist, or registered salesperson before using this medicine.**

- (1) Patients undergoing medical treatment from a doctor.
- (2) Pregnant women or women suspected of being pregnant.
- (3) Patients who have experienced allergic symptoms associated with drugs, etc.
- (4) Those with extensive lesions.
- (5) Those with severe moistening or erosion

**2. If the following symptoms appear after use, it may be a side effect, so stop using it immediately and consult your doctor, pharmacist, or registered salesperson with this case.**

Related area	Symptoms
Skin	Rash ▪ redness, itching, swelling, blotch, irritation, heat on skin, dryness, irritability
Skin (Affected area)	Ringworm such as athlete's foot and ringworm, acne, suppurative symptoms, and persistent irritation

**3. If symptoms do not improve after using between 5 and 6 days, stop using the medicine, consult a doctor, pharmacist, or registered salesperson registered seller with this case or this Japanese product manual. the Japanese instructions written on this package case, and consult a doctor, pharmacist or registered salesperson.**

### Directions

Apply an appropriate amount to the affected area several times a day.

### When using this product

- (1) This medicine should be given to children only under adult supervision.

- (2) Be careful not to get it in your eyes. If it gets into your eyes, immediately wash it with water or lukewarm water. If the symptoms are severe (such as persistent redness, pain, or unstoppable tearing), please consult an ophthalmologist.
- (3) This medicine is for external use only and should not be taken internally.

### Precautions for storage and handling

- (1) Store in a cool place free from direct sunlight by a seal tightly.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)
- (4) Do not touch your eyes or mucous membranes with hands covered with this medicine.
- (5) Be careful not to get it on glasses, watches, accessories, plastics, synthetic fiber products, leather products, painted surfaces such as floors or furniture, as it may cause deterioration.
- (6) Do not use this medicine past their expiration date. Also, please use it as soon as possible after opening, even if it is within the expiration date.

### Inactive ingredients:

Contains isopropyl palmitate, petrolatum, behenyl alcohol, stearyl alcohol, cetanol, glyceryl monostearate, cetomacrogol, parabens, dibutylhydroxytoluene, dimethylpolysiloxane, and diisopropanolamine.

### Contact information for the Relief System for Side Effects

Pharmaceuticals and Medical Devices Agency

[http://WWW.pmda.go.jp/kenkouhigai\\_cmp/index.html](http://WWW.pmda.go.jp/kenkouhigai_cmp/index.html)

Telephone: 0120-149-931 (toll-free)

### *Questions or comments?*

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.  
Customer consultation room ☎ 058-246-8771



Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

***[Request of limited correspondence]*** <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. ([www.kyowa-ci.co.jp](http://www.kyowa-ci.co.jp)).

***Disclaimer statement***

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.