

Product information

【Drug Facts】



＜Kyowa-Gifu-Japan＞

For resource saving the case is combined with the Japanese instruction written on the package case. Be sure to read the instruction carefully before use. Also, please keep it in a safe place so that you can read again, when you need it.

Class ② drugs **Skin Disease Treatment** (LIQUID)

＜Scalp Itching・Eczema＞

SHIN-TESITON EX GOLD EKI

- (1) 8 active ingredients are effective against severe scalp itching and eczema.
- (2) This medicine is blended with prednisolone valerate ester acetate (Antedrug).
- (3) It is comfortable to use with l-menthol and it is refreshing and refreshing.
- (4) It is a push-type dripping container that is easy to apply directly to the affected area.

Active ingredients (in 100mL)

Purpose

Prednisolone valeric acid ester acetic acid ester (Antedrug steroids PVA) 0.15g...	Reduce skin irritation
Diphenhydramine hydrochloride 2.0g....	Reduce the occurrence of itching
Crotamiton 3.0g.....	Reduce itching
Glycyrrhetic acid 0.2g.....	Reduce inflammation
l-Menthol 3.5g.....	Give a refreshing feeling
Lidocaine 1.0g.....	Reduce itching with local anesthesia
Isopropyl methyl phenol 0.1g.....	Bactericidal ingredient
Allantoin 0.2g.....	Remedies for skin

Uses

Itching, Eczema, Dermatitis, Rashes, Hives, Heat rashes,
Insect bites

Warnings

[To be avoided]

(Failure to observe the following may result in the current symptoms worsening or adverse reactions occurring.)

1. Do not use for the next part:

- (1) If you have the affected areas with chicken pox (water pox), Athlete's foot (Tinea pedis), Ringworm in the groin, etc. and/or who have purulent on the affected part.
- (2) Wound surface, around eyes, mucous membranes etc.

2. Please do not use it extensively on the face.

3. Do not use long-term continuously.

[Consultation]

1. The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

- (1) A person undergoing medical treatment from a doctor.
- (2) A pregnant woman or a woman suspected of being pregnant.
- (3) A person with a medical history of being allergic to medicines and other substances.
- (4) The affected area is spreaded on the wide.
- (5) A person with wet or awful sore on the affected part.

2. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, stop the use immediately and consult a doctor, pharmacist, or registered distributor with the Japanese instructions written on this package case.

Related area	Symptoms
Skin	Rash • redness, itching, blotch, swelling, irritation, heat on skin, dryness, irritability
Skin (Affected area)	Athlete's foot (Tinea pedis) • Ringworm in the groin by trichomonas, Acne, Purulent symptoms, persistent feeling of irritation

3. The following symptoms may appear after taking it, so if such symptoms persist or increase. Please stop taking it and bring the Japanese instructions written on the package case, consult a doctor, pharmacist or registered distributor.

Directions

Apply an appropriate amount to the affected area several times a day.

[When using this product]

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children only under adult supervision.
- (3) Be careful not to get into eyes. If it gets in your eyes, please wash immediately with water or lukewarm water. And If the symptoms are severe, please be given medical treatment by an eye doctor immediately.
- (4) Please use only for external treatment.
- (5) If the affected area has been become a wound, it may feel like sting at the time of application.

Other information

- (1) Store in a cool place free from direct sunlight by a seal tightly.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)
- (4) Do not get close to fire. In addition, please do not throw in fire after use.
- (5) Be careful not to adhere to metal objects such as glasses, watches and accessories, clothing, plastics, leather products, and painted surfaces such as floors and furniture, as they may deteriorate or discolor.
- (6) If it adheres to the dyed hair, it may be discolored.
- (7) Please close the cap tightly after use.
- (8) Do not use this medicine whose expiration date (described on the bottom of the box) has passed.

Inactive ingredients:

1,3-Butylene glycol, Macrogol, Ethanol,
Phosphoric acid, Hydrophobized
hydroxypropylmethylcellulose.

Questions or comments

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.
Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] **＜Kyowa-Gifu-Japan＞**

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the Revised date, the attached document is revised from time to time, so the English translation may not be the latest.