

# Product information

## 【Drug Facts】



<Kyowa-Gifu-Japan>

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 2 drugs

**Multivitamin Preparations (TABLETS)**

## SESHINRIKI U.P

“SESHINRIKI UP is a Vitamin-Amino acid containing health medicine which has the oxoamidine powder (garlic preparation with reduced odor), dried ginseng extract (root of ginseng root), dry extract of Epilepsy sutra (whole herb of ovary), dry extract of cashew (tuberous root of thrush kudami).

It is effective for nutritional support in the case of fever and debilitating diseases such as nourishment, weakness, physical fatigue and cold.

You can use it widely when you feel tired due to your daily work or stress, when you don't get tired of your body, or when you're concerned about your physical strength decline.

### *Active ingredients (in 4 tablets/daily dose)*

### *Purpose*

Oxoamidine powder 120mg .....	Herbal medicine
(equivalent to (12,000mg of the crude drug)	
L-Arginine hydrochloride 150mg .....	Amino acid supplementation
Ginseng dry extract 72mg.....	Antitussive expectorant
(equivalent to 720mg of the herbal medicine Ginseng Radix.)	
Epimedium Herb extract 60mg.....	Herbal medicine
(equivalent to 600mg of the crude drug)	
Polygonum Root extract 36mg .....	Herbal medicine
(equivalent to 360mg of the crude drug)	
Ursodeoxycholic acid 12mg .....	Bile promoter
Tocopherol succinate succinate,(Vitamin E) 20mg ....	Vitamin E supplementation
Gamma-Oryzanol 10mg .....	Gamma-Oryzanol supplementation

Thiamine nitrate 10mg .....

Riboflavin 6mg .....

Nicotinic acid amide 25mg .....

Anhydrous Caffeine 50mg .....

Vitamin B1  
supplementation  
Vitamin B2  
supplementation  
Nicotinamide  
supplementation  
Stimulant

## ***Uses***

- Nourishment.
- Weakness about physical.
- Nutritional supplementation in case of physical fatigue, after illness during illness, anorexia, malnutrition, fever wasting disease, pregnancy and nursing period.

## ***Warnings***

### ***[Consultation]***

- 1 The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

- (1) A person undergoing medical treatment from a doctor.
- (2) A person who have ever had allergic symptoms due to any medication.

2. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, stop the take immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

Related area	Symptoms
Skin	Rash · Redness、Itching
Digestive organ	Stomach discomfort

3. The following symptoms may appear after taking it, so if such symptoms persist or increase. Please stop taking it and consult a doctor, pharmacist or registered distributor with the Japanese package insert.

Diarrhea, Constipation.

4. If symptoms do not improve after taking this medicine for about one month, stop taking the medicine, bring the Japanese package insert, and consult a doctor, pharmacist or registered distributor.

5. After taking this medicine, menstruation may come earlier than planned, and menstrual blood volume may increase slightly. If the bleeding lasts for a long time, take this document and consult a doctor, pharmacist or registered distributor.

### ***Directions***

Please take the following amount after breakfast and dinner without chewing.

Age	One dose	Number of doses per day
adult and children 15 years and over	2 tablets	two times
children under 15 years	Do not use	

- (1) Please follow the recommended dosage and directions.
- (2) Don't worry about taking this medicine, if urine may become yellow, so this is due to riboflavin in this medicine.

### ***Other information***

- (1) Store in a cool and dry with less humidity place free from direct sunlight by a seal tightly.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (It may be misused or the quality may deteriorate.)
- (4) Do not touch the tablets with wet hands as may change its shape due to moisture.
- (5) The fillings in the bottle is to prevent the tablets from breaking down during shipping. It will be not necessary after opening the bottle, so please remove the fillings.
- (6) The "paper packet" in the bottle is a desiccant. Do not remove covered paper of it, and never take out, open and eat.
- (7) Please write the "Open Date" of this medicine on the inner lid of the case.
- (8) Do not take overdue products. In addition, please use within six months for quality maintenance after opening even if it is within expiration date.

***Inactive ingredients:***

cellulose, lactose, hydroxypropyl cellulose, caromelose Ca, sillic acid anhydride, stearic acidMg, hypromerolose, macrogol, vinyl acetate resin, polyvinylacetal diethylaminoacetate, calcium carbonate, talc, gum arabic, gelatin, polyoxyethylene polyoxypropylene glycol, sucrose, titanium oxide, yellow No. 5, carnauba wax.

***Questions or comments***

If you have any suggestions about this product, please contact the shop you purchased or the following address.

---

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.  
Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

***[Request of limited correspondence] <Kyowa-Gifu-Japan>***

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. ([www.kyowa-ci.co.jp](http://www.kyowa-ci.co.jp)).

***Disclaimer statement***

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the Revised date, the attached document is revised from time to time, so the English translation may not be the latest.